

Asthma Treatment Plan

(This asthma action plan meets NJ Law N.J.S.A. 18A:40-12.8) (Physician's Orders)

The Pediatric/Adult
Asthma Coalition
of New Jersey
"Your Pathway to Asthma Control"
Original PACNJ approved Plan available at
www.pacnj.org

Sponsored by
AMERICAN LUNG ASSOCIATION
of New Jersey



Please Print)

Name		Date of Birth	Effective Date
Doctor	Parent/Guardian (if applicable)		Emergency Contact
Phone	Phone		Phone

HEALTHY



You have **all** of these:

- Breathing is good
- No cough or wheeze
- Sleep through the night
- Can work, exercise, and play

And/or Peak flow above

Take daily medicine(s). All metered dose inhalers (MDI) to be used with spacers.

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
<input type="checkbox"/> Advair® <input type="checkbox"/> 100, <input type="checkbox"/> 250, <input type="checkbox"/> 5001 inhalation twice a day
<input type="checkbox"/> Advair® HFA <input type="checkbox"/> 45, <input type="checkbox"/> 115, <input type="checkbox"/> 2302 puffs MDI twice a day
<input type="checkbox"/> Asmanex® Twisthaler® <input type="checkbox"/> 110, <input type="checkbox"/> 220 <input type="checkbox"/> 1, <input type="checkbox"/> 2 inhalations a day
<input type="checkbox"/> Flovent® <input type="checkbox"/> 44, <input type="checkbox"/> 110, <input type="checkbox"/> 2202 inhalations twice a day
<input type="checkbox"/> Flovent® Diskus® 50 mcg1 inhalation twice a day
<input type="checkbox"/> Pulmicort Flexhaler® <input type="checkbox"/> 90, <input type="checkbox"/> 180 <input type="checkbox"/> 1, <input type="checkbox"/> 2 inhalations <input type="checkbox"/> once or <input type="checkbox"/> twice a day
<input type="checkbox"/> Pulmicort Respules® <input type="checkbox"/> 0.25, <input type="checkbox"/> 0.5, <input type="checkbox"/> 1.01 unit nebulized <input type="checkbox"/> once or <input type="checkbox"/> twice a day
<input type="checkbox"/> Qvar® <input type="checkbox"/> 40, <input type="checkbox"/> 802 inhalations twice a day
<input type="checkbox"/> Singulair <input type="checkbox"/> 4, <input type="checkbox"/> 5, <input type="checkbox"/> 10 mg1 tablet daily
<input type="checkbox"/> Symbicort® <input type="checkbox"/> 80, <input type="checkbox"/> 1602 puffs MDI twice a day
<input type="checkbox"/> Other	
<input type="checkbox"/> None	

Remember to rinse your mouth after taking inhaled medicine.

If exercise triggers your asthma, take this medicine _____ minutes before exercise.

Triggers

Check all items that trigger patient's asthma:

- Chalk dust
- Cigarette Smoke & second hand smoke
- Colds/Flu
- Dust mites, dust, stuffed animals, carpet
- Exercise
- Mold
- Ozone alert days
- Pests - rodents & cockroaches
- Pets - animal dander
- Plants, flowers, cut grass, pollen
- Strong odors, perfumes, cleaning products, scented products
- Sudden temperature change
- Wood Smoke
- Foods:

Other:

This asthma treatment plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs.

CAUTION



You have **any** of these:

- Exposure to known trigger
- Cough
- Mild wheeze
- Tight chest
- Coughing at night
- Other: _____

And/or Peak flow from _____ to _____

Continue daily medicine(s) and add fast-acting medicine(s).

MEDICINE	HOW MUCH to take and HOW OFTEN to take it
<input type="checkbox"/> Accuneb® <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Albuterol <input type="checkbox"/> 1.25, <input type="checkbox"/> 2.5 mg1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Albuterol <input type="checkbox"/> Pro-Air <input type="checkbox"/> Proventil®2 puffs MDI every 4 hours as needed
<input type="checkbox"/> Ventolin® <input type="checkbox"/> Maxair <input type="checkbox"/> Xopenex®2 puffs MDI every 4 hours as needed
<input type="checkbox"/> Xopenex® <input type="checkbox"/> 0.31, <input type="checkbox"/> 0.63, <input type="checkbox"/> 1.25 mg1 unit nebulized every 4 hours as needed
<input type="checkbox"/> Increase the dose of, or add:	

➡ If fast-acting medicine is needed more than 2 times a week, except before exercise, then call your doctor.

EMERGENCY



Your asthma is getting worse fast:

- Fast-acting medicine did not help within 15-20 minutes
- Breathing is hard and fast
- Nose opens wide
- Ribs show
- Trouble walking and talking
- Lips blue • Fingernails blue

And/or Peak flow below

Take these medicines NOW and call 911. Asthma can be a life-threatening illness. Do not wait!

- Accuneb® 0.63, 1.25 mg1 unit nebulized every 20 minutes
- Albuterol 1.25, 2.5 mg1 unit nebulized every 20 minutes
- Albuterol Pro-Air Proventil®2 puffs MDI every 20 minutes
- Ventolin® Maxair Xopenex®2 puffs MDI every 20 minutes
- Xopenex® 0.31, 0.63, 1.25 mg1 unit nebulized every 20 minutes
- Other

FOR MINORS ONLY:

- This student is capable and has been instructed in the proper method of self-administering of the inhaled medications named above in accordance with NJ Law.
- This student is not approved to self-medicate.

PHYSICIAN/APN/PA SIGNATURE _____ DATE _____

PARENT/GUARDIAN SIGNATURE _____

PHYSICIAN STAMP

Make a copy for patient and for physician file. For children under 18, send original to school nurse or child care provider.

The Pediatric/Adult Asthma Coalition of New Jersey, sponsored by the American Lung Association of New Jersey, and this publication are supported by a grant from the New Jersey Department of Health and Senior Services (NJHSS), with funding provided by the U.S. Centers for Disease Control and Prevention (CDC/DPPE) Cooperative Agreement #5U49CE000206-2. It contains no endorsement, approval, or disapproval of the views and do not necessarily represent the official views of NJHSS or the CDC/DPPE.

Although this document has been funded wholly or in part by the United States Environmental Protection Agency under Agreement Nos. 48054-01-4 and 48054-01-5 to the American Lung Association of New Jersey, it has not gone through the Agency's publication review process and therefore, may not necessarily reflect the views of the Agency and no official endorsement should be inferred.

EFFECTIVE MARCH 2008

Permission to reproduce blank form Approved by the New Jersey Thematic Society

**The Pediatric/Adult
Asthma Coalition
of New Jersey**

"Your Pathway to Asthma Control"
Original PACNJ approved Plan available at
www.pacnj.org

**Asthma Treatment Plan
Patient/Parent Instructions**



The PACNJ Asthma Treatment Plan is designed to help everyone understand the steps necessary for the individual patient to achieve the goal of controlled asthma.

1. Patients/Parents/Guardians: Before taking this form to your Health Care Provider:

Complete the top left section with:

- Patient's name
- Patient's date of birth
- Patient's doctor's name & phone number
- Parent/Guardian's name & phone number
- An Emergency Contact person's name & phone number

2. Your Health Care Provider will:

Complete the following areas:

- The effective date of this plan
- The medicine information for the Healthy, Caution and Emergency sections
- Your Health Care Provider will check the box next to the medication and circle how much and how often to take it
- Your Health Care Provider may check "OTHER" and:
 - ◆ Write in asthma medications not listed on the form
 - ◆ Write in additional medications that will control your asthma
 - ◆ Write in generic medications in place of the name brand on the form
- Together you and your Health Care Provider will decide what asthma treatment is best for you or your child to follow

3. Patients/Parents/Guardians & Health Care Providers together:

Discuss and then complete the following areas:

- Patient's peak flow range in the Healthy, Caution and Emergency sections on the left side of the form
- Patient's asthma triggers on the right side of the form
- For Minors Only section at the bottom of the form: Discuss your child's ability to self-administer the inhaled medications, check the appropriate box, and then both you and your Health Care Provider must sign and date the form

4. Parents/Guardians: After completing the form with your Health Care Provider:

- Make copies of the Asthma Treatment Plan and give the signed original to your child's school nurse or child care provider
- Keep a copy easily available at home to help manage your child's asthma
- Give copies of the Asthma Treatment Plan to everyone who provides care for your child, for example: babysitters, before/after school program staff, coaches, scout leaders

This Asthma Treatment Plan is meant to assist, not replace, the clinical decision-making required to meet individual patient needs. Not all asthma medications are listed and the generic names are not listed.

Disclaimers:

The use of this Website/PACNJ Asthma Treatment Plan and its content is at your own risk. The content is provided on an "as is" basis. The American Lung Association of the Mid-Atlantic (ALAM-A), the Pediatric/Adult Asthma Coalition of New Jersey and all affiliates disclaim all warranties, express or implied, statutory or otherwise, including but not limited to the implied warranties or merchantability, non-infringement of third parties' rights, and fitness for a particular purpose.

ALAM-A makes no representations or warranties about the accuracy, reliability, completeness, currency, or timeliness of the content. ALAM-A makes no warranty, representation or guaranty that the information will be uninterrupted or error free or that any defects can be corrected.

In no event shall ALAM-A be liable for any damages (including, without limitation, incidental and consequential damages, personal injury/wrongful death, lost profits, or damages resulting from data or business interruption) resulting from the use or inability to use the content of this Asthma Treatment Plan whether based on warranty, contract, tort or any other legal theory, and whether or not ALAM-A is advised of the possibility of such damages. ALAM-A and its affiliates are not liable for any claim, whatsoever, caused by your use or misuse of the Asthma Treatment Plan, nor of this website.

The Pediatric/Adult Asthma Coalition of New Jersey, sponsored by the American Lung Association of New Jersey, and this publication are supported by a grant from the New Jersey Department of Health and Senior Services (NJHSS), with funds provided by the U.S. Centers for Disease Control and Prevention (USCDCP) under Cooperative Agreement 5U59EH000205-2. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NJHSS or the USCDCP. Although this document has been funded wholly or in part by the United States Environmental Protection Agency under Agreements XA97256707-1, XA98284401-3 and XA97250908-0 to the American Lung Association of New Jersey, it has not gone through the Agency's publications review process and therefore, may not necessarily reflect the views of the Agency and no official endorsement should be inferred. Information in this publication is not intended to diagnose health problems or take the place of medical advice. For asthma or any medical condition, seek medical advice from your child's or your health care professional.

 **AMERICAN
LUNG
ASSOCIATION.**
of New Jersey